

DTIC FILE COPY

U

②

DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

AD-A198 376

2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA			1b. RESTRICTIVE MARKINGS NA		3. DISTRIBUTION/AVAILABILITY OF REPORT Distribution unlimited	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) 4B569			5. MONITORING ORGANIZATION REPORT NUMBER(S) NA			DTIC ELECTED AUG 26 1988 SE
6a. NAME OF PERFORMING ORGANIZATION University of Health Sciences/ The Chicago Medical School		6b. OFFICE SYMBOL (If applicable) NA	7a. NAME OF MONITORING ORGANIZATION Office of Naval Research			
6c. ADDRESS (City, State, and ZIP Code) 3333 Green Bay Road North Chicago, IL 60064		7b. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000				
8a. NAME OF FUNDING/SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable) ONR	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-86-K-0477			
8c. ADDRESS (City, State, and ZIP Code) 800 N. Quincy Street Arlington, VA 22217-5000			10. SOURCE OF FUNDING NUMBERS			
			PROGRAM ELEMENT NO. 61153N	PROJECT NO. PRO4106	TASK NO. NR4413201	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) Evolution of Phosphofructokinase						
12. PERSONAL AUTHOR(S) Robert G. Kemp						
13a. TYPE OF REPORT Annual		13b. TIME COVERED FROM 8/1/87 TO 7/31/88		14. DATE OF REPORT (Year, Month, Day) 7/31/88		15. PAGE COUNT 2
16. SUPPLEMENTARY NOTATION						
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)			
FIELD 08	GROUP	SUB-GROUP	protein evolution; protein engineering; directed mutagenesis; ()			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)						
<p>Comparative sequence data suggests that mammalian phosphofructokinase (PFK) has evolved from a procaryotic precursor by gene duplication, fusion, and mutation of previous catalytic sites into new regulatory ligand binding sites. We are examining this process by two approaches. First, we are attempting to duplicate these processes by recombinant DNA technology. The <i>E. coli</i> pfk gene is being recloned into a convenient plasmid. Using a synthetic oligonucleotide that matches the mammalian "link" region, two <i>E. coli</i> pfk genes will be joined. The properties of the expressed "duplex" enzyme will be analyzed. In the second approach, we are looking at several other PFKs to determine whether their evolution has followed a similar path. Those being examined include PPi dependent PFKs.</p>						
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS				21. ABSTRACT SECURITY CLASSIFICATION U		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. M. Marron				22b. TELEPHONE (Include Area Code) 202-696-4760		22c. OFFICE SYMBOL ONR

DD Form 1473, JUN 86

Previous editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

S/N 0102-LF-014-6603

88 8 25 195

PROJECT TITLE: EVOLUTION OF PHOSPHOFRUCTOKINASE

Annual Report for the period of 8/1/87 - 7/31/88

Principal Investigator: Robert G. Kemp, Ph.D.

Department of Biological Chemistry & Structure
University of Health Sciences/
The Chicago Medical School
3333 Green Bay Road
North Chicago, IL 60064

(1)

Work on the construction of the fused duplex PFK gene progressed to the point of complete assembly of the link between two *E. coli* PFK genes. Insertion of the link into the appropriate position has been deferred until additional, more convenient constructs of the gene are prepared, which is part of the second area of effort. Our initial construct, pSLJ3, was inconvenient for studies of site-directed mutagenesis because of difficulty in moving back and forth to M13. Our initial altered construct that was to solve this problem was reported last year but proved to be unsuccessful due to instability. We have now solved the problem by using a Bal digestion through the strong promoter that interfered with M 13 expression, readdition of linkers, and insertion into both pUC plasmids or M13.

(2)

We have characterized further our first site-directed mutant of *E. coli* PFK. A mutation was made by the method of Mandecki, which involves introducing a single restriction site cleavage into the gene and transforming with the cut plasmid plus an oligonucleotide that bridges the cut and contains one or more mismatches near the restriction site. We cleaved pSLJ3 with BglII and added a 65-mer that removed the restriction site and changed G174 to E. is three residues from critical catalytic residue involving ATP and seems to be important for the folding of the pocket. The kinetic properties of the mutant enzyme are fairly similar to native enzyme except for a low V_{max} . Further studies have shown that the low V_{max} was apparent and that the actual reason for the low values was the instability of the enzyme. Heat denaturation studies demonstrated that the mutant was denatured rapidly at 40 to 42°C whereas the wild-type enzyme denatured in the range of 60 to 62°C.

(3)

We extended our interest in the evolution of the PFKs to an evaluation of the place in the scheme of the PPi-dependent PFK of plants. A procedure was developed for the purification of inorganic pyrophosphate; fructose-6-phosphate 1-phospho-transferase (PPi-PFK) from potato tubers. The enzyme has the structure $\alpha_4\beta_4$ with a subunit of 68kDa and a β subunit of 60 kDa. The structural relationship of the enzyme to other PFKs and to fructose

bisphosphatase was examined by immunoprecipitation and immunoblotting. Antibodies to the plant enzyme did not react with *E. coli* PFK. No crossreaction was seen among the following enzymes or their antibodies: yeast fructose bisphosphatase; rabbit PFKs A,B, or the enzyme from brain; and the two subunits of the potato P_{Pi}-PFK. On the other hand, antibody to *E. coli* PFK-1 strongly cross-reacts with the 60 kDa polypeptide but not 68 kDa peptide.

The data support the idea that the plant P_{Pi}-PFK has evolved from the same procaryotic progenitor that led to the major ATP-dependent enzyme of *E. coli* and to the mammalian PF-1-Ks. The structural identity of the second subunit, α , and its role in the reaction remain to be established. Three short tryptic peptides of the 60 kDal and one from the 68 kDal have been sequenced. Oligonucleotides are being synthesized on the basis of these sequences to probe a plant cDNA library.

In addition, we have recently partially purified the ATP dependent PFK of potato tubers. Using the Western blot technique, we have shown that the antibody to the 60 kDal subunit of the P_{Pi}-PFK is capable of recognizing a subunit of the ATP dependent enzyme.

Accession For	
NTIS	GRA&I <input checked="" type="checkbox"/>
DTIC	TAB <input type="checkbox"/>
Unannounced <input type="checkbox"/>	
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

